

AD 646047

YALE UNIVERSITY
Department of Statistics

Technical Report No. 1
November, 1966

"A geometrical approach to the special stable distributions"

By L. J. SAVAGE

This research was supported by the Army, Navy, Air Force and NASA under a contract administered by the Office of Naval Research. Reproduction in whole or in part is permitted for any purpose of the United States Government.


Contract number: Nonr 609(52)

Task number: NR 042-242

FACILITY FORM 602

N67-2617.1

(ACCESSION NUMBER)		(THRU)	
9		0	
(PAGES)		(CODE)	
ad 646047		19	
(NASA CR OR TMX OR AD NUMBER)		(CATEGORY)	
CR 83499			



Acquisitioned Document
90T

The convolution formulas for the Cauchy distribution and for the distribution of the reciprocal of the square of a standard normal random variable are here derived from the geometry of circularly symmetric distributions on the plane. Other known derivations seem quite different, and in a sense, less elementary, though some are by no means less interesting. A few other illustrations of the geometrical method are also included.

Let a , a' , b , and b' be points on a circle with $a \neq a'$ and $b \neq b'$, and suppose that the chords (a, a') and (b, b') intersect in a point c not exterior to the circle and not the same as a or b . Then the angle (a, c, b) is half the sum of the arcs (a, b) and (a', b') . This fact seems old and widely known. The special case in which $a' = b' = c$ is very familiar indeed (Euclid III 26 and 27), and the general case is easily inferred from the special one with the aid of the construction line (a, b') .

If a and b are two points in the Euclidean plane, the direction from a to b is the vector from a to b normalized to unit length, and the unoriented direction is the unordered couple consisting of the direction and its negative, so that the unoriented direction from a to b is also the unoriented direction from b to a . The notion of a uniformly distributed random direction needs no explicit definition here, and a random unoriented direction will be called uniformly distributed if it has the same distribution as the unoriented direction associated with a uniformly distributed random direction. In this terminology, the geometric fact of the preceding paragraph has the following probabilistic interpretation.

Lemma 1. If a random point P is uniformly distributed on the periphery of a circle in the plane and c is not exterior to that circle, then the unoriented direction from c to P is uniformly distributed.

For the present form of the lemma and of the argument leading to it, I thank E. J. G. Pitman and Paul Lévy, respectively.

Let Z be a random vector distributed with circular symmetry in the Euclidean vector plane. Assume, to avoid unimportant complications, that $\Pr(Z = 0) = 0$. Let X and Y be a pair of cartesian coordinates of Z , and let R and A be the corresponding polar coordinates, so that:
 $X = R \cos A$; $Y = R \sin A$; R and A are independent; $\Pr(R > 0) = 1$; and A is uniformly distributed on the reals modulo 2π .

For this note, interest centers on the example in which X and Y are independent and normally distributed with mean 0 and variance 1, which will be called the standard example. The more general situation is treated mainly to emphasize the structure of the arguments.

Let " \sim " mean "is distributed like".

Theorem 1. For all real f and g , $fX + gY \sim (f^2 + g^2)^{1/2} X$.

Proof: $fX + gY = (f^2 + g^2)^{1/2} R \cos(A - \alpha)$, where $\alpha = \arctan f/g$. But $A - \alpha \sim A$ and is independent of R . Since $X = R \cos A$, this completes the proof.

Specialized to the standard example, Theorem 1 is the convolution formula for the normal distribution. The proof, however familiar, helps to set the stage.

Lemma 2. If σ and τ are nonnegative and not both 0,

$$\frac{XY}{(\sigma^2 X^2 + \tau^2 Y^2)^{1/2}} \sim \frac{X}{\sigma + \tau}.$$

Proof: Expressed in polar coordinates, the lemma says that R times a certain function of A is distributed like $R \cos A$ or, equivalently $R \sin A$. Since R and A are independent, it will be adequate to show that the function of A , namely

$$h(A) = \frac{(\sigma + \tau) \sin A \cos A}{(\sigma^2 \cos^2 A + \tau^2 \sin^2 A)^{1/2}},$$

is distributed like $\sin A$. But

$$h(A/2) = \frac{(\sigma + \tau) \sin A}{\{2(\sigma^2 + \tau^2) + 2(\sigma^2 - \tau^2) \cos A\}^{1/2}}$$

$$= \frac{\sin A}{\{1 + 2\rho \cos A + \rho^2\}^{1/2}},$$

where, $\rho = (\sigma - \tau)/(\sigma + \tau)$. Since $(\sin 2A, \cos 2A) \sim (\sin A, \cos A)$, what needs to be shown is that $h(A/2) \sim \sin A$ for each ρ in $[-1, 1]$.

Geometrically, if A is thought of as a random point P on the unit circle, then $h(A/2)$ is the sine of the angular coordinate B of that point as viewed from the point $c = (-\rho, 0)$. If the angular coordinate associated with the direction from c to P is B , that associated with the opposite direction is $B + \pi$, the sine of which is the negative of that of B . Therefore, $|\sin B|$ depends only on the unoriented direction from c to P . But according to Lemma 1, this unoriented direction is distributed just as it would be if B were uniformly distributed, that is, $|\sin B| \sim |\sin A|$. This, together with the remark that the distribution of $\sin A$ and $\sin B$ are both symmetric, completes the proof of the present lemma.

Corollary 1. If M and N are independent and normal with mean 0, then $L = MN/(M^2 + N^2)^{1/2}$ is normal with mean 0 and $s.d.(L) = (s.d.(M)^{-1} + s.d.(N)^{-1})^{-1}$.

This corollary has been proved (Shepp 1964) by recognizing its obvious equivalence to the standard-example application of the next theorem, which itself is obviously equivalent to the lemma.

Theorem 2. For nonnegative p and q ,

$$\frac{p}{X^2} + \frac{q}{Y^2} \sim \frac{(p^{1/2} + q^{1/2})^2}{X^2}$$

Applied to the standard example, the theorem asserts that the distribution of the reciprocal of the square of a standard normal variable is stable of order $1/2$. Doetsch (1936) attributes the fact to Cesaro and gives two

interesting, but not very probabilistic, demonstrations. Lévy (1940) independently discovered the fact in the course of a beautiful investigation of the Wiener process. For some further information, pursue "Stable distributions of order 1/2" in the Index of (Feller 1966).

The standard Cauchy distribution is the distribution of $C = Y/X = \tan A$.

Theorem 3. For all real f and g not both 0,

$$(fC + g)^{-1} \sim \frac{fC + g}{(f^2 + g^2)}.$$

Proof:

$$\begin{aligned} (fC + g)^{-1} &= \frac{X}{fY + gX} \\ &= \frac{f}{(f^2 + g^2)} \frac{(-gY + fX)}{(fY + gX)} + \frac{g}{f^2 + g^2} \\ &\sim \frac{f}{(f^2 + g^2)} \frac{Y}{X} + \frac{g}{f^2 + g^2}, \end{aligned}$$

as in the proof of Theorem 1.

Theorem 3 has been proved (Mann 1962, p. 1270) by direct calculation with the Cauchy density.

Theorem 4. If $Z' = (X', Y')$ is independent of Z and it too is distributed with circular symmetry, then, for all real f and g ,

$$(1) \quad f \frac{X'}{X} + g \frac{Y'}{Y} \sim (|f| + |g|) \frac{X'}{X}.$$

Proof: The left side of (1) is U/V , where

$$U = \frac{fYX' + gXY'}{(g^2X^2 + f^2Y^2)^{1/2}}$$

and

$$V = \frac{XY}{(g^2X^2 + f^2Y^2)^{1/2}}.$$

Given X and Y , U is distributed like X' , as Theorem 1 shows. Therefore U is independent of V and distributed like X' . According to Lemma 2, $V \sim X/(|f| + |g|)$. This proves the theorem.

Corollary 2. If C and C' are independent and have the standard Cauchy distribution, then for all f and g , $fC + gC' \sim (|f| + |g|)C$.

This popular elementary fact has been demonstrated in several other ways. See for example (Feller 1966, p. 50). The proof given here is easily extended to the multivariate Cauchy distribution, that is, the distribution of a normally distributed vector divided by an independent, standard normally distributed number.

REFERENCES

- Gustav Doetsch, Thetarelationen als Konsequenzen des Huygensschen und Eulerischen Prinzips in der Theorie der Wärmeleitung. *Mathematische Zeitschrift*, 40, (1936) 613-628.
- William Feller, An Introduction to Probability Theory and Its Applications: Volume II, (1966). John Wiley and Sons, New York.
- Paul Lévy, Sur certains processus stochastiques homogènes. *Compositio Mathematica*, 7, (1940) 283-339.
- M. V. Menon, a Characterization of the Cauchy Distribution. *Annals of Mathematical Statistics*, 33, (1962) 1267-1271.
- Larry Shepp, Problem 62-9, Normal Functions of normal random variables. *SIAM Review*, 6, (1964) 459-460.

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Yale University		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE A geometrical approach to the special stable distributions.			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical Report, November 1966.			
5. AUTHOR(S) (Last name, first name, initial) Savage, Leonard J.			
6. REPORT DATE November 1966		7a. TOTAL NO. OF PAGES 6	7b. NO. OF REFS 5
8a. CONTRACT OR GRANT NO. Nonr 609(52) a. PROJECT NO. RR 003 05 01 c. Task No. NR 042-242 d.		9a. ORIGINATOR'S REPORT NUMBER(S) Technical Report No. 1 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Logistics and Mathematical Statistics Branch, Office of Naval Research, Washington, D.C. 20360	
13. ABSTRACT If X and Y are independent and normal about 0 , possibly with different variances, then $Z = XY/(X^2 + Y^2)^{1/2}$ is also normally distributed about 0 . This is known and known to be tantamount to the stability of the distribution of X^{-2} . The present paper reduces the distribution of Z to a very elementary fact of plane geometry and shows geometrically that it also implies the stability of the Cauchy distribution.			

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Stable law Normal distribution Cauchy distribution Geometrical approach in probability						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.